

NAPLEX^{Q&As}

North American Pharmacist Licensure Examination

Pass Test Prep NAPLEX Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.leads4pass.com/naplex.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers



QUESTION 1

How many kcal per gram does IV dextrose provide?

- A. 0.9 kcal/g
- B. 1.2 kcal/g
- C. 1.5 kcal/g
- D. 3.4 kcal/g

Correct Answer: D

IV dextrose provides 3.4 kcal/g.

QUESTION 2

A patient who weighs 80kg is ordered Esmolol at 50mcg/kg/min. Esmolol comes in 2500mg/250 ml NS premixed bags. What is the infusion rate in mls/hr?

- A. 20mls/hr
- B. 6mls/hr
- C. 8mls/hr
- D. 24mls/hr
- E. 32mls/hr

Correct Answer: D

QUESTION 3

What is the weight of 1000 ml of serum protein whose specific gravity is 1.27?

- A. 1270 gm
- B. 127mg
- C. 12.7mg
- D. 12.7gm
- E. 1.27gm

Correct Answer: A

SG= weight/mL, $1.27 = x/1000\text{ml}$, $x = 1270\text{gm}$

QUESTION 4

If you mix 30 gm 5% lidocaine cream and 90gm of 0.5% hydrocortisone cream, what percent of lidocaine and hydrocortisone do you have as the end product?

- A. Lidocaine/Hydrocortisone 2%/1.25%
- B. Lidocaine/Hydrocortisone 0.375%/0.15%
- C. Lidocaine/Hydrocortisone 1.25 %/ 0.15%
- D. Lidocaine/Hydrocortisone 2% /0.25%
- E. Lidocaine/Hydrocortisone 1.25% /0.375%

Correct Answer: E

QUESTION 5

Which of the following Anti-epileptic medication can cause pancreatitis?

- A. Carbamazepine
- B. Gabapentin
- C. Valproic acid
- D. Levetiracetam
- E. Phenobarbital

Correct Answer: C

Cases of life-threatening pancreatitis have been reported in both pediatric and adult patients receiving valproic acid or its analogs. Patients should be warned that abdominal pain, nausea, vomiting, and/or anorexia can be symptoms of pancreatitis that require prompt medical evaluation. If pancreatitis is diagnosed, valproate should be discontinued.

Reference: <http://www.clinicalpharmacology-ip.com/Forms/Monograph/monograph.aspx?cpnum=637andsec=moncontrandt=0>

[Latest NAPLEX Dumps](#)

[NAPLEX VCE Dumps](#)

[NAPLEX Braindumps](#)