

NCLEX-RN^{Q&As}

National Council Licensure Examination(NCLEX-RN)

Pass NCLEX NCLEX-RN Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.leads4pass.com/nclex-rn.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by NCLEX
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers



QUESTION 1

As the nurse assesses a male adolescent with chlamydia, the nurse determines that a sign of chlamydia is:

- A. Enlarged penis
- B. Secondary lymphadenitis
- C. Epididymitis
- D. Hepatomegaly

Correct Answer: C

(A)

An enlarged penis is not a sign of chlamydia. (B) Secondary lymphadenitis is a complication of lymphogranuloma venereum. (C) Untreated chlamydial infection can spread from the urethra, causing epididymitis, which presents as a tender, scrotal swelling.

(D)

Hepatomegaly is not a complication.

QUESTION 2

A 35-year-old client is admitted to the hospital for elective tubal ligation. While the nurse is doing preoperative teaching, the client says, "The anesthesiologist said she was going to give me balanced anesthesia. What exactly is that?" The best explanation for the nurse to give the client would be that balanced anesthesia:

- A. Is a type of regional anesthesia
- B. Uses equal amounts of inhalation agents and liquid agents
- C. Does not depress the central nervous system
- D. Is a combination of several anesthetic agents or drugs producing a smooth induction and minimal complications

Correct Answer: D

(A) Regional anesthesia does not produce loss of consciousness and is indicated for excision of moles, cysts, and endoscopic surgeries. (B) Varying amounts of anesthetic agents are used when employing balanced anesthesia. Amounts depend on age, weight, condition of the client, and surgical procedure. (C) General anesthesia is a drug-induced depression of the central nervous system that produces loss of consciousness and decreased muscle activity. (D) Balanced anesthesia is a combination of a number of anesthetic agents that produce a smooth induction, appropriate depth of anesthesia, and appropriate muscle relaxation with minimal complications.

QUESTION 3

A 5-year-old child cries continually in her bed. Her parents have been unsuccessful in assisting her in expressing her feelings. Which activity should the nurse provide the child to assist her in expressing her feelings?

- A. Books with colorful pictures
- B. Music
- C. Riding toys
- D. Puppets

Correct Answer: D

(A) Books increase cognition, assist with fine motor skills, and augment language development. (B) Music provides auditory stimulation and large-muscle activity. (C) Riding toys provide large-muscle activity. (D) Puppets allow expression of feelings and fears that otherwise could not be directly communicated.

QUESTION 4

An elective saline abortion has been performed on a 3- week primigravida. Following the procedure, the nurse should be alert for which early side effect?

- A. Water satiety
- B. Thirst
- C. Edema
- D. Diabetes insipidus

Correct Answer: B

(A) If the client is experiencing water satiety, there is no more desire for water. (B) Absorption of saline into circulation rather than into amniotic sac increases serum sodium and desire for water. (C) Edema can be a late side effect caused by water intoxication. (D) Diabetes insipidus occurs as a result of deficient antidiuretic hormone.

QUESTION 5

The physician prescribes a medical regimen of isoniazid, rifampin, and vitamin B6 for a tuberculosis client. The nurse instructs the client that B6 is given because it:

- A. Increases activity of isoniazid
- B. Increases activity of rifampin
- C. Improves nutritional status
- D. Reduces peripheral neuropathy

Correct Answer: D

(A) Vitamin B6 does not enhance the activity of isoniazid. (B) Vitamin B6 does not enhance the activity of rifampin. (C) A vitamin alone does not improve nutritional status. (D) Isoniazid leads to Vitamin B6 deficiency, which is manifested as peripheral neuropathy.

[NCLEX-RN Practice Test](#)

[NCLEX-RN Study Guide](#)

[NCLEX-RN Braindumps](#)